

Minimal diagrams of virtual links: II

Vassily Olegovich Manturov

Abstract

In the present paper we bring together minimality conditions proposed in papers [10, 11] and present some new minimality conditions for classical and virtual knots and links.

1 The main result

This paper is a sequel of [10, 11]. We deal with virtual link diagrams and test whether these diagrams are minimal with respect to the number of classical crossings. All necessary definitions can be found in [6, 10, 11].

Throughout the text, all virtual links are thought to be non-split, unless otherwise specified. In any minimality theorem for links we assume that the link in question has no split diagrams. All virtual links are thought to be orientable in the sense of atoms, see [6, 10].

First, let us formulate the two main theorems from [10] and [11].

The main Theorem from [10] says the following:

Theorem 1. *If a virtual link diagram L' is 1-complete and 2-complete then it is minimal.*

The main Theorem from [11] says the following:

Theorem 2. *Suppose the diagram K of a classical knot is good. Then it is minimal in the classical category. In other words, if the diagram K has n crossings then for any classical diagram representing the same knot the number of crossing is at least n .*

In the sequel, we shall refer to these two theorems as “The first Theorem” and “The second Theorem”. The first theorem says that if the span of the Kauffman polynomial for a virtual link diagram is “as large as it should be” and the genus of the corresponding atom is “as large as it should be” then the diagram is minimal.

The second theorem deals only with classical knots (not virtual knots, and not links) and it has only one condition. This condition says that no cell of the corresponding atom touches itself at a crossing. This condition is much stronger than the first minimality condition used in the first theorem saying that *the leading and the lowest term of the Kauffman state sum expansion do not vanish*. The condition of the second Theorem says that each of the two extreme coefficients equals precisely to one, and we have only one non-trivial summand contributing to each of them.

In fact, the condition of the second Theorem allows to consider the cablings of the initial diagram L , the diagrams $D_m(L)$ where m is a positive

integer. It turns out that the main condition of the second Theorem (that the diagram is good) is hereditary: if it holds for L , then it should hold for $D_m(L)$. This allows to establish minimality by passing to cablings and some more tricks (see [11]). Here we do not worry about the thickness of the Khovanov homology [12] of the corresponding atom.

The first condition of the first Theorem (saying that the leading and the lowest terms in the Kauffman state-sum expansion) are **not** hereditary: they may hold for L , but not for $D_m(L)$, $m \neq 1$, and usually they **do not**. For instance, they do not hold in the classical case. Thus, to establish the minimality of a (virtual) knot or link diagram we have to handle the genus thus adding one more condition on the Khovanov homology.

In the present paper, we wish to formulate a stronger statement explaining the connection between the first Theorem and the second Theorem, namely, we shall deal with the genus of links $D_m(L)$, and $D_m(L\#\bar{L})$, where $\bar{L}$ is the mirror image of the diagram L .

First, we deal with classical knots (to have the connected sum operation well defined). At the end of the present paper, we shall prove a theorem on virtual links.

If some auxiliary theorem or lemma admits a formulation for virtual links, we formulate it for the general case though we might need it only for the case of classical knots.

Suppose we have a virtual link diagram L with n classical crossings such that the corresponding atom has genus g and Euler characteristic $\chi = 2 - 2g$. Then the maximal possible span for the Kauffman bracket of L is estimated as ([10]):

$$span\langle L \rangle \leq 4n + 2(\chi - 2). \quad (1)$$

Also, the Khovanov homology has thickness $T(L)$ defined by the diagonals it lives between, see [11].

In [11] (using a refined version of the result from [13]) we demonstrated that

$$T(L) \leq 2 + g(L). \quad (2)$$

We shall refer to the corresponding equalities as “the span of the Kauffman bracket is as large as it should be” and “the Khovanov homology is as thick as it should be”.

Let K be a classical knot diagram with n crossings, $N = 2n$. The usual estimate for $span\langle D_m \rangle$ is $2(m^2 + m)N + 2m\chi(K\#\bar{K}) - 4$. It is so if the neither the leading term nor the lowest term of the Kauffman state-sum expansion vanishes. The following asymptotic theorem says that if this length is smaller then expected, but asymptotically the difference between the real length and the estimate is not very large, then the initial diagram is minimal. Namely, we have

Theorem 3 (The asymptotic theorem). *Suppose that for some $\varepsilon > 0$ there is an infinite sequence $i_1 < i_2 < \dots i_m < \dots$ of positive integers such that for any positive integer m we have*

$$span\langle D_{i_m}(K\#\bar{K}) \rangle \geq 2(i_m^2 + i_m)N + 2i_m\chi(K\#\bar{K}) - 4 - (4 - \varepsilon)(i_m^2 + i_m).$$

Then the diagram K is minimal in the classical category.

We have the following

Lemma 1. *If a virtual link diagram L is good then $T(L) \geq g(L)$.*

Proof. Indeed, one should just take the A -state with v_- associated to all circles and the B -state with v_+ associated to all circles. The property that the diagram is good guarantees that the first chain is a cycle, whence the second one is not a boundary. Recalling the definition of the atom genus, we get the required estimate. $\square$

Now, having a good diagram L of a classical knot, we see that all diagrams $D_m(L \# \bar{L})$ are also good. By Lemma 1 we see that the diagram L obviously satisfies the condition of Theorem 3. Here we may take a constant for ζ_m . Thus, the diagram L is minimal.

So, the estimate for the thickness of the Khovanov homology for a good diagram L is that it is in between $2 + g(L)$ and $g(L)$.

This immediately results in the following

Theorem 4. *Let L be a good virtual link diagram with n classical crossings. Then any virtual link diagram L' equivalent to L has at least $n - 2$ classical crossings.*

Proof. Indeed, suppose we have a diagram L' equivalent to L with n' classical crossings. Then the thickness of L' is at least $g(L)$, so the genus $g(L')$ is at least $g(L) - 2$, thus, $\chi(L') \leq \chi(L) + 4$. From this we see that

$$4n + 2(\chi(L) - 2) = \text{span}\langle L \rangle = \text{span}\langle L' \rangle \leq 4n' + 2(\chi(L) + 2), \quad (3)$$

so $4(n - n') \leq 8$, that completes the proof. $\square$

We are still unable to use the trick with connected summation with mirror image as in [11] and prove the exact result (in the unframed category). However, the reasonings with Khovanov homology give a minimality estimate between $n - 2$ and n classical crossings.

Proof of Theorem 3. The proof goes in the same lines as that of the second Theorem [10].

Indeed, fix a positive integer m . Suppose there is a classical diagram K' having n' crossings ($n' < n$) and representing the same classical knot as K .

The diagrams $D_{i_m}(K \# \bar{K})$ and $D_{i_m}(K' \# \bar{K}')$ generate isotopic knots. Denote $D_{i_m}(K \# \bar{K})$ by D_m and denote $D_{i_m}(K' \# \bar{K}')$ by D'_m . By definition we have $\langle D_m \rangle = \langle D'_m \rangle$. Also, set $\chi = \chi(K \# \bar{K})$, $\chi' = \chi(K' \# \bar{K}')$.

We have:

$$\text{span}\langle D_m \rangle \geq 4i_m^2 N + 2(\chi_m - 2) - (4 - \varepsilon)(i_m^2 + i_m), \quad (4)$$

where $\chi_m = \chi(D_{i_m})$. The atom $V(D_m)$ has $i_m^2 N$ vertices, $2i_m^2 N$ edges and $i_m \Gamma$ 2-cells, where $\Gamma = N + \chi$ is the number of the 2-cells of the atom $K \# \bar{K}$. Thus,

$$\text{span}\langle D_{i_m} \rangle \geq 2(i_m^2 + i_m)N + 2i_m \chi - 4 - (4 - \varepsilon)(i_m^2 + i_m). \quad (5)$$

For D'_m we have:

$$\text{span}\langle D'_m \rangle \leq 2(i_m^2 + i_m)N + 2i_m\chi' - 4. \quad (6)$$

Thus, taking into account $\langle D'_m \rangle = \langle D_m \rangle$, we get

$$2(i_m^2 + i_m)(N - N') \leq (4 - \varepsilon)(i_m^2 + i_m) + 2i_m(\chi' - \chi). \quad (7)$$

This leads to a contradiction since $N - N' \geq 2$. $\square$

Remark 1. *If we deal with classical link diagrams then for any link diagram L which is not good and for any positive integer $m > 1$, the leading coefficient in the Kauffman state-sum expansion for $\langle D_m(L) \rangle$ is equal to zero. Thus, we can not apply Theorem to the classical case directly.*

In the virtual case, there are some diagrams L which are not good, but for every $m > 1$, the state-sum expansion of the Kauffman polynomial for $D_m(L)$ does not vanish. But here we can not apply Theorem 1 directly because the connected summation for virtual diagrams is not well defined.

Besides the asymptotic theorem (which works in the case of classical knots only, because it uses the connected sum construction), we also have the following generalisation of the First theorem:

Remark 2. *Theorem 3 works also in the case of long virtual knots. The proof is literally the same because we have a well-defined connect summation for such knots.*

In the long virtual category, we have a lot of examples where the first minimality condition is hereditary. Such examples obviously give us minimal diagrams by theorem 3.

Also, analogous theorems remain true for tangles and braids. We shall discuss it in separate papers.

References

- [1] Kauffman, L.H. (1987), State Models and the Jones Polynomial, *Topology*, **26** (1987), pp. 395–407.
- [2] Kauffman, L. H. (1999), Virtual knot theory, *European Journal of Combinatorics* **20**(7), pp. 662–690.
- [3] Jones, V. F. R. (1985), A polynomial invariant for links via Neumann algebras, *Bull. Amer. Math. Soc.*, **129**, pp. 103–112.
- [4] Khovanov, M. (1997), A categorification of the Jones polynomial, *Duke Math. J.*, **101** (3), pp.359-426.
- [5] Kuperberg, G. (2002), What is a Virtual Link?, www.arXiv.org, math-GT/0208039, *Algebraic and Geometric Topology*, 2003, **3**, 587–591.
- [6] Manturov, V.O., Teoriya uzlov, *Regular and Chaotic Dynamics*, Moscow-Izhevsk, 2005.
- [7] Manturov V.O. (2003), Kauffman-like polynomial and curves in 2-surfaces, *Journal of Knot Theory and Its Ramifications*, **12**, (8), pp.1145-1153.

- [8] Manturov, V.O. (2004), Long virtual knots and their invariants, *Journal of Knot Theory and Its Ramifications*, **13** (8), pp. 1029-1039
- [9] Manturov, V.O., The Khovanov Complex for Virtual Links, Arxiv:GT/ 0501317.
- [10] Manturov, V.O., Minimal diagrams of classical and virtual links, ArXiv:GT/ 0501393.
- [11] Manturov, V.O., Minimal diagrams of classical knots, ArXiv:GT/ 0501510.
- [12] Shumakovitch, A., Torsion of the Khovanov homology, Arxiv:GT/ 0405474.
- [13] Wehrli, S., A spanning tree model for the Khovanov homology, Arxiv: GT/ 0409328